

Claims:

1. A method of vaccinating a mammal against a disease state, comprising administering to said mammal, within an appropriate vector, a nucleotide sequence encoding an antigenic peptide associated with the disease state;

additionally administering to said mammal a compound which enhances both humoral and cellular immune responses initiated by the antigenic peptide, the compound being selected from:

- 4-(2-formyl-3-hydroxyphenoxy)methyl)benzoic acid;
5-(2-formyl-3-hydroxyphenoxy)pentanamide;
N,N-diethyl 5-(2-formyl-3-hydroxyphenoxy)pentanamide;
N-isopropyl 5-(2-formyl-3-hydroxyphenoxy)pentanamide;
ethyl 5-(2-formyl-3-hydroxyphenoxy)pentanoate;
5-(2-formyl-3-hydroxyphenoxy)pentanonitrile;
(±)-5-(2-formyl-3-hydroxyphenoxy)-2-methylpentanoic acid;
5-(2-formyl-3-hydroxyphenoxy)-2,2-dimethylpentanoic acid;
methyl 3-(2-formyl-3-hydroxyphenoxy)methylbenzoate;
3-(2-formyl-3-hydroxyphenoxy)methylbenzoic acid;
benzyl 5-(2-formyl-3-hydroxyphenoxy)pentanoate;
5-[4-(2-formyl-3-hydroxyphenoxy)-N-butyl]tetrazole;
7-(2-formyl-3-hydroxyphenoxy)heptanoic acid;
5-(2-formyl-3-hydroxy-4-n-propoxyphenoxy)pentanoic acid;
5-(4,6-dichloro-2-formyl-3-hydroxyphenoxy)pentanoic acid;
5-(2-formyl-3-hydroxyphenoxy)-N-methylsulphonylpentanamide;
ethyl 4-(2-formyl-3-hydroxyphenoxy)methyl)benzoate;
5-(4-chloro-2-formyl-3-hydroxyphenoxy)pentanoic acid;
5-(3-acetylamino-2-formylphenoxy)pentanoic acid;
Aminoguanidine;
4-(2-formyl-3-hydroxyphenoxy)butanoic acid;
6-(2-formyl-3-hydroxyphenoxy)hexanoic acid;
ethyl 4-(3-acetylamino-2-formylphenoxy)methyl)benzoate;
4-(3-acetylamino-2-formylphenoxy)methyl)benzoic acid;
2-(2-formyl-3-hydroxyphenoxy)methyl)benzoic acid;

5- [4-(2-formyl-3-hydroxyphenoxy)methyl]phenyl]tetrazole;
5-(2-formyl-3-hydroxy-4-methoxyphenoxy)pentanoic acid;
3-(2-formyl-3-hydroxyphenoxy)propionitrile;
4-Hydroxyphenylacetaldehyde;
5 Phenylacetaldehyde;
4-Methoxyphenylacetaldehyde;
1-hydroxy-2-phenylpropane;
3-Phenylpropanaldehyde;
4-Nitrobenzaldehyde;
Methyl 4-formylbenzoate;
4-Chlorobenzaldehyde;
4-Methoxybenzaldehyde;
4-Methylbenzaldehyde;
8,10-Dioxoundecanoic acid;
15 4,6-Dioxoheptanoic acid;
Pentanedione;
5-methoxy-1-tetralone;
6-methoxy-1-tetralone;
7-methoxy-1-tetralone;
20 2-tetralone;
3-hydroxy-1-(4-methoxyphenyl)-3-methyl-2-butanone;
2',4'-dihydroxy-2-(4-methoxyphenyl)acetophenone;
2-hydroxy-1-(4-methoxyphenyl)-pent-2-ene-4-one;
Naringenin 4',5,6-trihydroxyflavonone;
25 4'-methoxy-2-(4-methoxyphenyl)acetophenone;
6,7-dihydroxycoumarin;
7-methoxy-2-tetralone;
6,7-dimethoxy-2-tetralone;
6-hydroxy-4-methylcoumarin;
30 Homogentisic acid gamma lactone;
6-hydroxy-1,2-naphthoquinone;
8-methoxy-2-tetralone;

and physiologically acceptable salts thereof, where appropriate.

SUB 10
B2
C01N 7/1
TOTO 2916

Sub D1

2. The method according to claim 1 wherein administration of the compound takes place on between one and seven occasions, between about 14 days prior to and about 14 days post administration of the nucleotide sequence.

5 3. The method according to claim 1 wherein administration of the compound takes place on between one and seven occasions, between about 7 days prior to and about 7 days post administration of the nucleotide sequence.

10 4. The method according to claim 1 wherein administration of the compound takes place between about 24 hours prior to and about 24 hours post administration of the nucleotide sequence.

15 5. The method according to claim 1 wherein administration of the compound is substantially simultaneous with administration of the nucleotide sequence.

A 20 6. The method according to ~~any one of claims 1 to 5~~ ^{claim 1} which is repeated between 1 and 4 times, at intervals of between about 1 day and about 18 months.

A 25 7. The method according to ~~any one of claims 1 to 6~~ ^{claim 1} wherein administration of the nucleotide sequence is via the oral, nasal, pulmonary, intramuscular, subcutaneous or intradermal routes.

30 8. The method according to claim 7 wherein the nucleotide sequence is administered using a gene-gun delivery technique.

A Sub D2 9. The method according to ~~any one of claims 1 to 8~~ ^{claim 1} wherein administration of the compound is via the oral, nasal, pulmonary, intramuscular, subcutaneous, intradermal or topical routes.

A 10. The method according to ~~any one of claims 1 to 8~~ ^{claim 1} wherein the compound is administered using a gene-gun delivery technique.

A. Sub
DB

11. The method according to ~~either claim 9 or claim 10~~ wherein the compound is administered at a dose of between about 0.1mg and about 100 mg/per kg per administration.

Claim 9

A 5

12. The method according to ~~any one of claims 1 to 11~~ wherein the mammal is a human.

claim 1

A

10

13. The method according to ~~any one of claims 1 to 12~~ wherein the compound is 4-(2-formyl-3-hydroxyphenoxy)methylbenzoic acid.

claim 1

15

14. A vaccine composition comprising a nucleotide sequence which encodes for an antigenic peptide associated with a disease state and which is within an appropriate vector, and a compound which will enhance both humoral and cellular immune responses in a mammal which are initiated by the antigenic peptide, the compound being selected from:

- 4-(2-formyl-3-hydroxyphenoxy)methylbenzoic acid;
- 5-(2-formyl-3-hydroxyphenoxy)pentanamide;
- N,N-diethyl 5-(2-formyl-3-hydroxyphenoxy)pentanamide;
- N-isopropyl 5-(2-formyl-3-hydroxyphenoxy)pentanamide;
- ethyl 5-(2-formyl-3-hydroxyphenoxy)pentanoate;
- 5-(2-formyl-3-hydroxyphenoxy)pentanonitrile;
- (±)-5-(2-formyl-3-hydroxyphenoxy)-2-methylpentanoic acid;
- 5-(2-formyl-3-hydroxyphenoxy)-2,2-dimethylpentanoic acid;
- methyl 3-(2-formyl-3-hydroxyphenoxy)methylbenzoate;
- 3-(2-formyl-3-hydroxyphenoxy)methylbenzoic acid;
- benzyl 5-(2-formyl-3-hydroxyphenoxy)pentanoate;
- 5-[4-(2-formyl-3-hydroxyphenoxy)-N-butyl]tetrazole;
- 7-(2-formyl-3-hydroxyphenoxy)heptanoic acid;
- 5-(2-formyl-3-hydroxy-4-n-propoxyphenoxy)pentanoic acid;
- 5-(4,6-dichloro-2-formyl-3-hydroxyphenoxy)pentanoic acid;
- 5-(2-formyl-3-hydroxyphenoxy)-N-methylsulphonylpentanamide;
- ethyl 4-(2-formyl-3-hydroxyphenoxy)methylbenzoate;
- 5-(4-chloro-2-formyl-3-hydroxyphenoxy)pentanoic acid;
- 5-(3-acetylamino-2-formylphenoxy)pentanoic acid;

25

30

35

09763463
2949260
SUB
C1

5
SUB
C1
CON 10
09763462.050.001
15
20
25
30
35

Aminoguanidine;
4-(2-formyl-3-hydroxyphenoxy)butanoic acid;
6-(2-formyl-3-hydroxyphenoxy)hexanoic acid;
ethyl 4-(3-acetylamino-2-formylphenoxy)methyl)benzoate;
4-(3-acetylamino-2-formylphenoxy)methyl)benzoic acid;
2-(2-formyl-3-hydroxyphenoxy)methyl)benzoic acid;
5-[4-(2-formyl-3-hydroxyphenoxy)methyl)phenyl]tetrazole;
5-(2-formyl-3-hydroxy-4-methoxyphenoxy)pentanoic acid;
3-(2-formyl-3-hydroxyphenoxy)propionitrile;
4-Hydroxyphenylacetaldehyde;
Phenylacetaldehyde;
4-Methoxyphenylacetaldehyde;
1-hydroxy-2-phenylpropane;
3-Phenylpropanaldehyde;
4-Nitrobenzaldehyde;
Methyl 4-formylbenzoate;
4-Chlorobenzaldehyde;
4-Methoxybenzaldehyde;
4-Methylbenzaldehyde;
8,10-Dioxoundecanoic acid;
4,6-Dioxoheptanoic acid;
Pentanedione;
5-methoxy-1-tetralone;
6-methoxy-1-tetralone;
7-methoxy-1-tetralone;
2-tetralone;
3-hydroxy-1-(4-methoxyphenyl)-3-methyl-2-butanone;
2',4'-dihydroxy-2-(4-methoxyphenyl)acetophenone;
2-hydroxy-1-(4-methoxyphenyl)-pent-2-en-4-one;
Naringenin 4',5,6-trihydroxyflavonone;
4'-methoxy-2-(4-methoxyphenyl)acetophenone;
6,7-dihydroxycoumarin;
7-methoxy-2-tetralone;
6,7-dimethoxy-2-tetralone;
6-hydroxy-4-methylcoumarin;

SUB
C1
CONT.

Homogentisic acid gamma lactone;
6-hydroxy-1,2-naphthoquinone;
8-methoxy-2-tetralone;

5 namely and physiologically acceptable salts thereof, where appropriate.

15. The vaccine composition according to claim 14 which is in a form suitable for administration via the oral, nasal, pulmonary, intramuscular, subcutaneous or intradermal route.

10 16. The vaccine composition according to claim 14 which is in a form suitable for administration using a gene-gun delivery technique.

15 17. The vaccine composition according to ~~any one of claims 14 to 16~~ ^{claim 14} wherein the compound is 4-(2-formyl-3-hydroxyphenoxy)methylbenzoic acid.

20 18. Use of a compound in the manufacture of a medicament, wherein administration of the compound to a mammal enhances both humoral and cellular responses initiated by an antigenic peptide associated with a disease state, peptide being expressed as a result of administration to said mammal of a nucleotide sequence encoding for the peptide;

wherein said compound is selected from:

SUB
CA

25 4-(2-formyl-3-hydroxyphenoxy)methylbenzoic acid;
5-(2-formyl-3-hydroxyphenoxy)pentanamide;
N,N-diethyl 5-(2-formyl-3-hydroxyphenoxy)pentanamide;
N-isopropyl 5-(2-formyl-3-hydroxyphenoxy)pentanamide;
ethyl 5-(2-formyl-3-hydroxyphenoxy)pentanoate;
30 5-(2-formyl-3-hydroxyphenoxy)pentanonitrile;
(±)-5-(2-formyl-3-hydroxyphenoxy)-2-methylpentanoic acid;
5-(2-formyl-3-hydroxyphenoxy)-2,2-dimethylpentanoic acid;
methyl 3-(2-formyl-3-hydroxyphenoxy)methylbenzoate;
3-(2-formyl-3-hydroxyphenoxy)methylbenzoic acid;
35 benzyl 5-(2-formyl-3-hydroxyphenoxy)pentanoate;

09763462.050101

- 5- [4-(2-formyl-3-hydroxyphenoxy)-N-butyl]tetrazole;
7-(2-formyl-3-hydroxyphenoxy)heptanoic acid;
5-(2-formyl-3-hydroxy-4-n-propoxyphenoxy)pentanoic acid;
5-(4,6-dichloro-2-formyl-3-hydroxyphenoxy)pentanoic acid;
5- (2-formyl-3-hydroxyphenoxy)-N-methylsulphonylpentanamide;
ethyl 4-(2-formyl-3-hydroxyphenoxy)methyl)benzoate;
5-(4-chloro-2-formyl-3-hydroxyphenoxy)pentanoic acid;
5-(3-acetylamino-2-formylphenoxy)pentanoic acid;
Aminoguanidine;
4-(2-formyl-3-hydroxyphenoxy)butanoic acid;
6-(2-formyl-3-hydroxyphenoxy)hexanoic acid;
ethyl 4-(3-acetylamino-2-formylphenoxy)methyl)benzoate;
4-(3-acetylamino-2-formylphenoxy)methyl)benzoic acid;
2-(2-formyl-3-hydroxyphenoxy)methyl)benzoic acid;
5-[4-(2-formyl-3-hydroxyphenoxy)methyl)phenyl]tetrazole;
5-(2-formyl-3-hydroxy-4-methoxyphenoxy)pentanoic acid;
3-(2-formyl-3-hydroxyphenoxy)propionitrile;
4-Hydroxyphenylacetaldehyde;
Phenylacetaldehyde;
4-Methoxyphenylacetaldehyde;
1-hydroxy-2-phenylpropane;
3-Phenylpropanaldehyde;
4-Nitrobenzaldehyde;
Methyl 4-formylbenzoate;
4-Chlorobenzaldehyde;
4-Methoxybenzaldehyde;
4-Methylbenzaldehyde;
8,10-Dioxoundecanoic acid;
4,6-Dioxoheptanoic acid;
Pentanedione;
5-methoxy-1-tetralone;
6-methoxy-1-tetralone;
7-methoxy-1-tetralone;
2-tetralone;
3-hydroxy-1-(4-methoxyphenyl)-3-methyl-2-butanone;

sub
C2
C6H5

10763462-050101

2',4'-dihydroxy-2-(4-methoxyphenyl)acetophenone;
 2-hydroxy-1-(4-methoxyphenyl)-pent-2-ene-4-one;
 Naringenin 4',5,6-trihydroxyflavonone;
 4'-methoxy-2-(4-methoxyphenyl)acetophenone;
 6,7-dihydroxycoumarin;
 7-methoxy-2-tetralone;
 6,7-dimethoxy-2-tetralone;
 6-hydroxy-4-methylcoumarin;
 Homogentisic acid gamma lactone;
 6-hydroxy-1,2-naphthoquinone;
 8-methoxy-2-tetralone;

and physiologically acceptable salts thereof, where appropriate.

19. The use according to claim 18 wherein the medicament is in a form suitable for administration via the oral, nasal, pulmonary, intramuscular, subcutaneous or intradermal routes.

20. The use according to claim 19 wherein the medicament is in a form suitable for administration using a gene-gun delivery technique.

21. The use according to ^{claim 18} ~~any one of claims 18 to 20~~ wherein the compound is 4-(2-formyl-3-hydroxyphenoxymethyl)benzoic acid.

22. The use according to ^{claim 18} ~~any one of claims 18 to 21~~ wherein the compound is administered at a dose of between about 0.1 mg/kg and 100 mg/kg per administration.

23. The use according to ^{claim 18} ~~any one of claims 18 to 22~~ wherein the medicament further comprises the nucleotide sequence.

24. A combination of components for separate, sequential or concomitant administration in a method according to claim 1, comprising the nucleotide sequence encoding an antigenic peptide and the compound which enhances both cellular and humoral immune responses initiated by the antigenic peptide.

SUB
C2
CDP

09763462-050101

A

A

A

SUB
C3